Adult Environmental Emergencies: Animal Bites and Stings

Snake Bites

I. All Provider Levels

- 1. Refer to the Patient Care Protocols.
- 2 Place the patient supine.



Note Well: Do not pack in ice.

Do not make any incisions.

- 3. Locate the fang mark(s), clean and treat the open wound.
- 4. Remove any rings, bracelets or other constricting items on the bitten extremity.
- 5. Immobilize any bitten extremities.
- 6. If evacuation will be prolonged, assess the patient's urine for signs of myoglobinuria (tea-like discoloration).
- 7. If myoglobinuria is found, treat with IV fluids.



Note Well: If possible, capture or identify the animal.

II. The Following Options are Available by Medical Control Only

- 1. Apply cool pack to the affected area.
- 2. Apply light constricting bands.
 - A. Each band should be about two inches from the wound.
 - B. Never place the bands on either side of a joint.

Effective Date: 1 May 2002 Revision Number: N/A

Revision Date: N/A Page E5.1

Adult Environmental Emergencies: Animal Bites and Stings

Animal Bites

I. All Provider Levels

- 1. Refer to the Patient Care Protocols.
- 2. Irrigate the wound.
- 3. Dress wound.



Note Well: Notify MPD and/or the Animal Control Office. If possible, identify, or capture the animal for rabies testing.

Bee Stings

I. All Provider Levels

- 1. Refer to the Patient Care Protocols.
- 2. Apply cool pack to reduce pain.
- 3. Monitor for signs of anaphylaxis and if present, refer to the Anaphylaxis/Allergic Reaction Protocol, D2.

Effective Date: 1 May 2002 Revision Number: N/A
Revision Date: N/A Page E5.2